

Metric Prefix

A metric prefix is a unit prefix that precedes a basic unit of measure to indicate a multiple or fraction of the unit.

In the previous section, we saw that Mhz means a million Hertz. Here's a list of the most common ones:

Name	Symbol	Base 10	Decimal
Tera	T	10^{12}	1 000 000 000 000
Giga	G	10^9	1 000 000 000
Mega	M	10^6	1 000 000
Kilo	k	10^3	1 000
hecto	h	10^2	100
deca	da	10^1	10
		10^0	1
deci	d	10^{-1}	0.1
centi	c	10^{-2}	0.01
milli	m	10^{-3}	0.001
micro	μ	10^{-6}	0.000 001
nano	n	10^{-9}	0.000 000 001
pico	p	10^{-12}	0.000 000 000 001

The decibel

A decibel (dB) is a way of saying how many times bigger (or smaller) something is compared to something else. For example, let's use the faintest sound that the human ear can detect as the basis. A normal conversation would be a million times louder, and a jack hammer would be a hundred billion times louder. To represent these vast differences in numbers, we use a logarithmic scale based on powers of 10.

Here's the idea. Start with representing numbers by exponents of 10:

Number	Base 10	Exponent
1 =	10^0	0
10 =	10^1	1
100 =	10^2	2
1000 =	10^3	3
10 000 =	10^4	4
100 000 =	10^5	5
1 000 000 =	10^6	6

If your math is a little rusty, notice how the exponent counts the number of zeros after the one (1 000 000 has 6 zeros so it's equal to 10^6)

For now, let's call the exponent a "bel" (not decibel yet). For example:

- Instead of saying that a sound is **1000 times** louder, we could say that it's **3 bels** louder, and
- Instead of saying that it's **10,000 times** louder, we could say that it's **4 bels** louder.

So the bel is just the number of zeros after the 1. But what if the number doesn't start with 1. For example: what if it's 5000 times louder?

- Since 5000 is between 1000 and 10,000, it might be tempting to say that it would 3.5 bels louder (midway between 3 and 4), but let's check to see if it works: $10^{3.5} = 3162$, which is between 1000 and 10,000, but not 5000.
- By trial and error, you might get pretty close: $10^{3.7} = 5011...$ so **5000** times louder is roughly **3.7 bels**.
- But a faster way of finding the exponent of a base 10 number is to use the **Log** button on your calculator: $\log(5000) \approx 3.699$ ¹⁾

In the same way that *milli* means a thousandth, and *centi* means a hundredth (think millimetre and centimetre), a *deci* means a tenth. So 1 bel contains 10 decibels (like 1 metre contains 10 decimetres, 100 centimetres, or 1000 millimetres).

So another way of saying 3.7 bels is to say 37 decibels, or 37 dB. So here's our table again with a few common “in between” numbers:

Number	Base 10	Exponent (bels)	decibels (dB)
1 =	10^0	0	0
2 ≈	$10^{0.3}$	0.3	3
4 ≈	$10^{0.6}$	0.6	6
8 ≈	$10^{0.9}$	0.9	9
10 =	10^1	1	10
100 =	10^2	2	20
1000 =	10^3	3	30
10 000 =	10^4	4	40
100 000 =	10^5	5	50
1 000 000 =	10^6	6	60

In addition to being more convenient to represent big numbers, decibels also allow us to **multiply** big (or small) numbers more easily by **adding** the dB instead. For example, look at:

- $100 \times 1000 = 100,000$

Let's write the same thing again in powers of 10:

- $10^2 \times 10^3 = 10^5$

Notice that while the numbers are multiplying (100×1000), the exponents are adding ($2 + 3$). So 100×1000 is the same as saying $20 \text{ dB} + 30 \text{ dB} = 50 \text{ dB}$. That is, adding the dB representation of numbers, is the same as multiplying those numbers.

Look back at the previous table; if you double the power, three times in a row:

- The final power is $2 \times 2 \times 2 = 8$ times more powerful than the original.
- In terms of dB, we can instead add 3dB three times: $3\text{dB} + 3\text{dB} + 3\text{dB} = 9\text{dB}$.

Here's a quick exercise: Using only the table above, what is the dB representation of 4000?

A 4000 fold increase is the same as a 1000 fold followed by two doubling. That is:

- $1000 \times 2 \times 2 = 4000$

Which means:

- $30\text{dB} + 3\text{dB} + 3\text{dB} = 36\text{dB}$


Quick calculator check: $10^{3.6} = 3981 \approx 4000$

So the only thing to memorize with decibels is that:

- a bel (10 dB) is the number of zeros after the 1. So that takes care of knowing 10dB, 20dB, 30dB, ...
- 3dB is one doubling.

With these two things, we can now estimate a bunch of in between numbers like 36dB (30dB + 3dB + 3dB), or 13dB (10dB + 3dB)

Here's Dave explaining this in more details:



Alternative Formulation

If you like formulas, what I did above was define the dB *implicitly* as:

$$\text{ratio} = 10^{\left(\frac{\text{dB}}{10}\right)}$$

For example, a ratio of 1000 is equivalent to 30 dB because:

$$1000 = 10^{\left(\frac{\text{30}}{10}\right)}$$

But what you'll often find in books is the following *explicit* definition:

$$\text{dB} = 10 \log \left(\text{ratio} \right)$$

While the two definitions are mathematically equivalent²⁾, I personally prefer the *implicit* definition because I find it easier to picture the dB in terms of powers of 10, where as I can't really picture the log function that well. But they are equivalent, so use the one you like most.

dBm

A related measurement is the dBm. While the decibel (dB) is a ratio between two quantities (saying “20 dB” is the same as saying “100 times more”), the dBm is a ratio between one quantity and 1 mW. That is, it's a measure of how much stronger (or weaker) the power of something is compared to 1 milliwatt.

For example, a typical fibre optic light signal to a house for internet has a light level of -16 dBm. Before we go through the math, recall that 16 dB = 10 dB + 3 dB + 3 dB. Which means, the ratio is $10 \times 2 \times 2 = 40$. Let's now see why -16 dBm is equal to a power of 25 μ W (microwatts):

$$-16 \text{ dBm} = 1 \text{ mW} \div 10^{1.6} \approx 1 \text{ mW} \div 40 = 0.025 \text{ mW} = 25 \mu\text{W}$$

Notice how the minus sign in front of the dBm means that the power is less (not more) so it divides (not multiplies).

Binary Numbers

Base 10

It's kind of weird explaining how to count in base 10 because we're so used to it. But to understand other bases, it helps to understand how base 10 really works first.

So base 10 means we have 10 different symbols to represent numbers, and when we need more, we just add more digits to the number. For example, let's count and pay close attention to what we're really doing:

- With one digit, we can count to nine because we have ten symbols: 0, 1, 2, 3, 4, 5, 6, 7, 8, 9
- To continue past nine, we need to add a new digit in front of the “counting digit”, and then we just do what we did above again: 10, 11, 12, 13, 14, 15, 16, 17, 18, 19
- And that new digit will increase in the same way until we run out of symbols: 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, \vdots 90, 91, 92, 93, 94, 95, 96, 97, 98, 99

- At which point we need to add a third digit (and so on).

It's strange to think about this but the ability to count is completely separate from having a sense of how big the numbers are. What I mean is: if you know how to change the symbols (from 0 to 1, and 1 to 2, and 2 to 3, etc) and how to add new digits (9 to 10, 99 to 100, ...) you can count forever without having a “number sense”. In base 10, it's hard to imagine not knowing how big a number is, but we'll actually do this in base 2 in a moment. You'll see, it's a weird feeling!

But first, here's another weird way to think of numbers. Take 234 for example. Let's “decompose” it into its powers of 10:

$$\begin{aligned} 234 &= 2 \times 100 + 3 \times 10 + 4 \times 1 \\ &= 2 \times 10^2 + 3 \times 10^1 + 4 \times 10^0 \end{aligned}$$

This is a very tedious way of expressing a simple number, but I promise it'll be invaluable when we try to convert from base 2 to base 10 in a moment (can you think of why?).

Base 2

Now let's do the same thing again in base 2. Imagine we only have two symbols we can use to count: 0 and 1. Let's see what counting to fifteen looks like starting at zero:

0, 1, 10, 11, 100, 101, 110, 111, 1000, 1001, 1010, 1011, 1100, 1101, 1110, 1111

Do you see the pattern? Here's another way of writing it with leading zeros and keeping track of where we are in base 10:

Base 2	Base 10
0000	0
0001	1
0010	2
0011	3
0100	4
0101	5
0110	6
0111	7
1000	8
1001	9
1010	10
1011	11
1100	12
1101	13
1110	14
1111	15

You see what I mean about being able to count without knowing how big the number is? All we're doing is “flipping bits” from 0 to 1 in the right order. If we wanted, we could count to 1001010011010 and have no idea of how big

that number is unless we kept track of it in base 10. That's because over the years, we developed a number sense for base 10 numbers, but we didn't for base 2 numbers since we don't typically use them.

Conversion

Now let's look at how to convert from one base to the other without counting.

Base 2 to Base 10

We'll use our example of 1101 in base 2 since we already know it's 13 in base 10. Before we do this, however, let's use subscripts to denote the base. For example, we'd write:

$$1101_2 = 13_{10}$$

Now let's see how we convert from base 2 to base 10 without counting. In the same way that each position in base ten is a factor of 10, each position in base 2 is a factor of 2:

$$\begin{aligned} 1101_2 &= 1 \times 2^3 + 1 \times 2^2 + 0 \times 2^1 + 1 \times 2^0 \\ &= 1 \times 8 + 1 \times 4 + 0 \times 2 + 1 \times 1 \\ &= 8 + 4 + 0 + 1 \\ &= 13_{10} \end{aligned}$$

Let's try that again with the crazy number I mentioned earlier: 1001010011010. Skipping the powers with a coefficient of 0 (to save space):

$$\begin{aligned} 1001010011010_2 &= 2^{12} + 2^9 + 2^7 + 2^4 + 2^3 + 2^1 \\ &= 4096 + 512 + 128 + 16 + 8 + 2 \\ &= 4762_{10} \end{aligned}$$

Base 10 to Base 2

Converting the other way around is a bit different, but kind of makes sense if you think about it. Here are the steps to convert 13_{10} back to 1101_2 and I'll explain a bit more after:

Division by 2	In Decimal	As fraction	Quotient	Remainder
$13 \div 2$	6.5	$6 + \frac{1}{2}$	6	1
$6 \div 2$	3.0	$3 + \frac{0}{2}$	3	0
$3 \div 2$	1.5	$1 + \frac{1}{2}$	1	1
$1 \div 2$	0.5	$0 + \frac{1}{2}$	0	1

Now we read the remainders from the bottom up: 1101.

Recall that the remainder is what's left of a division; it's not the decimal portion. For example:

$$22 \div 5 = 4.4 = 4 \frac{2}{5} = 4 \text{ remainder } 2 \text{ (of } 5)$$

In our case, the remainder is what's left after we divided by 2. So why does this method work?

What we really want to do is decompose 13 into the sum of its powers of 2. That is:

$$13_{10} = 8 + 4 + 0 + 1 = 2^3 + 2^2 + 0 + 2^0 = (1 \text{ quad } 1 \text{ quad } 0 \text{ quad } 1)_2$$

When we have a non-zero power of 2, we represent that position with a 1, when we have a zero power of 2, we write 0.

What the table above is doing is systematically dividing by 2 to see if the result is a whole number or not. If it is, it means that that position is 0 because the factor of two can be absorbed further up the chain. For example, the fact that $13 \div 2 = 4.5$ means that the first digit has to be 1.

Let's look at a trivial example converting 2037 from base 10 to base 10 (trivial indeed!):

Division by 10	In Decimal	As fraction	Quotient	Remainder
$2037 \div 10$	203.7	$203 + \frac{7}{10}$	203	7
$203 \div 10$	20.3	$20 + \frac{3}{10}$	20	3
$20 \div 10$	2	$2 + \frac{0}{10}$	2	0
$2 \div 10$	0.2	$0 + \frac{2}{10}$	0	2

Other Bases?

There are other useful bases such as hexadecimal (base 16). Here's an example of counting to twenty in hex:

0, 1, 2, 3, 4, 5, 6, 7, 8, 9, A, B, C, D, E, F, 10, 11, 12, 13, 14

The convention to denote a hexadecimal number is to use either 14_{16} or $0x14$.

The same logic applies to convert between hexadecimal and base 10:

$$\begin{aligned} \text{F61B}_{16} &= 15 \times 16^3 + 6 \times 16^2 + 1 \times 16^1 + 11 \times 16^0 \\ &= 15 \times 4096 + 6 \times 256 + 1 \times 16 + 11 \\ &= 63003_{10} \end{aligned}$$

One reason why hexadecimal is so useful in computer science is because one hexadecimal “digit” can represent four binary “digits”. For example,

Decimal	Hexadecimal	Binary
1	1	1
2	2	10
3	3	11
4	4	100
...		
7	7	111
8	8	1000
...		
15	F	1111
16	10	1 0000
...		
31	1F	1 1111
32	20	10 0000
...		
63	3F	11 1111
64	40	100 0000
...		
127	7F	111 1111
128	80	1000 0000

Decimal	Hexadecimal	Binary
...		
255	FF	1111 1111
256	100	1 0000 0000
...		
4095	FFF	1111 1111 1111
4096	1000	1 0000 0000 0000



1)

Depending on your calculator, you might have to type it in this order **5Log**

2)

These two definitions are equivalent because the log function is the inverse of the power function. That is, it helps us find the exponent when we know the other quantities. Mathematically, we say that: $a = 10^b \quad \Leftrightarrow \quad b = \log_{10}(a)$