

Optional Math of Waves

This is a brief survey of the math required to analyze waves at the first or second year university level. If you did well in grade 12 high school math, you'll probably be able to follow this and learn some new and really cool math.

Real numbers

If $(5)^2 = 25$ and $(-5)^2 = 25$, what number can you put in the box so that:

$$\Box^2 = -25$$

It turns out that there is no *real* number such that when you multiply it by itself you get a negative number. But could we invent an *imaginary* one?

Complex Numbers

Let's create an *imaginary* number called i such that:

$$i = \sqrt{-1} \quad \Rightarrow \quad i^2 = -1$$

Even though i is nowhere on the real line (in math, we say that: $i \notin \mathbb{R}$), we can none-the-less perform interesting mathematical operations with it:

- We can add it to a real number and create a *complex* number:

$$(1 + i)$$

- We can multiply complex numbers together:

$$(1+i)^2 = 1 + 2i + i^2 = 1 + 2i - 1 = 2i$$

- We can find roots:

$$z^4 = 16 \Rightarrow z^2 = \left\{ \begin{array}{l} 4 \\ -4 \end{array} \right\} \Rightarrow z = \pm 2 \text{ or } \pm 2i$$

A Little Philosophy

If these weird numbers follow all of the algebra rules without inconsistencies, does it mean they *exist* as much as the real numbers? Aren't complex numbers a mere *creation* by mad mathematicians? How about mathematics itself: is it *discovered* or *invented*?¹⁾

In a certain way, negative numbers are just as weird as complex numbers: after all, we know what 5 cars look like,

but what does -5 cars mean? And yet, in certain context (like temperature), we have no problem using negative numbers. Could it be that there are contexts where complex numbers make sense?

The Complex Plane

In the same way that we can represent real numbers by a point on the real number line...²⁾:



... we can also represent a complex number graphically on a complex plane by using the horizontal axis for the real part and the vertical axis for the imaginary part. For example, $(1 + i)$ would be represented as a point 45° up the horizontal axis and $\sqrt{2}$ away from the origin:

Download [polar.ggb](#)

You can move the point around to look at other complex numbers on the plane.

To convert between the Cartesian (a, b) and the Polar $(r \angle \theta)$ representations, only simple trigonometry and Pythagoras is needed.

$(a, b) \rightarrow r \angle \theta$	$r \angle \theta \rightarrow a + ib$
$r^2 = a^2 + b^2$	$a = r \cos \theta$
$\tan \theta = \frac{b}{a}$	$b = r \sin \theta$

Note that very often, we use radians instead of degrees for the angle. There are a total 360° or 2π radians in a circle. While most people are used to degrees, a radian is actually much easier to picture:

- Imagine a circle.
- Now imagine the length from the centre to the circle (along the —radius—).
- Take that length and lay it down on the perimeter of the circle.
- The angle that this length covers is 1 radian (because of the length of the radius on the circle).
- That's why a circle has 2π (because the circumference is $2\pi r$)

Roots

The complex plane has many useful applications, but one of them allows us to visualize roots of the form $z^n = w$. For example, if we set $w = 9$ and $n = 2$ on the graph below, we'll see that the roots of $z^2 = 9$ are $z = \pm 3$.

Download [complexroots.ggb](#)

Without using the graph above, what do you expect the solution(s) to $z^3 = 8$ will be? That is, what number(s), when multiplied by itself three times gives 8?

Now move $w = 8$ and $n = 3$ to have a look at the solutions graphically, you might be surprised by what you find.



So $z = 2$ was to be expected since $2^3 = 8$ but it looks like there's two more solutions.

To find them, we first notice that the three solutions are spread out evenly around the circle, that is they are 120° apart. So in polar coordinates, the three solutions are $z = 2\angle 0^\circ$, $2\angle 120^\circ$, $2\angle 240^\circ$.

We can now convert them to Cartesian:

- $z = 2\angle 0^\circ = (2\cos(0^\circ), 2\sin(0^\circ)) = (2, 0) = 2$
- $z = 2\angle 120^\circ = (2\cos(120^\circ), 2\sin(120^\circ)) = (-1, \sqrt{3}) = -1 + i\sqrt{3}$
- $z = 2\angle 240^\circ = (2\cos(240^\circ), 2\sin(240^\circ)) = (-1, -\sqrt{3}) = -1 - i\sqrt{3}$

Let's check that the second solution works:
$$\begin{aligned} (-1 + i\sqrt{3})^3 &= (-1 + i\sqrt{3}) \cdot (-1 + i\sqrt{3}) \cdot (-1 + i\sqrt{3}) \\ &= (-1 + i\sqrt{3}) \cdot (1 - 2i\sqrt{3} + (i^2)(3)) \\ &= (-1 + i\sqrt{3}) \cdot (1 - 2i\sqrt{3} - 3) \\ &= (-1 + i\sqrt{3}) \cdot (-2 - 2i\sqrt{3}) \\ &= 2 + 2i\sqrt{3} - 2i\sqrt{3} - 2(i^2)(3) \\ &= 2 - 2(-1)(3) \\ &= 2 + 6 = 8 \end{aligned}$$

The Euler Identity

The Euler identity exposes a deep relationship between trigonometric and exponential functions³⁾:

$$e^{i\theta} = \cos \theta + i \sin \theta$$

Let's use two different ways to verify that this mysterious identity is true.

The Derivatives

If we separate this identity into two functions and take their derivatives, we notice that:
$$\begin{aligned} f(\theta) &= e^{i\theta} & g(\theta) &= \cos \theta + i \sin \theta \\ f'(\theta) &= i e^{i\theta} & g'(\theta) &= -\sin \theta + i \cos \theta \end{aligned} \implies f'(\theta) = i \cdot f(\theta) \text{ and } g'(\theta) = i \cdot g(\theta)$$

We know that there's only one function $h(x)$ that satisfies the differential equation $h'(x) = ah(x)$, and it is $h(x) = A e^{ax}$. What Euler discovered is that when $a = i$, there's a second function that also satisfies the same differential equation! These two functions must therefore be one and the same.

Taylor

Another method to verify the Euler identity is to use Taylor series:

$$\begin{aligned} e^x &= 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \frac{x^5}{5!} + \cdots \\ \sin x &= x - \frac{x^3}{3!} + \frac{x^5}{5!} - \cdots \\ \cos x &= 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \cdots \end{aligned}$$

1. Replace x with $i\theta$ in the Taylor series of e^x
2. Replace x with θ in the Taylor series for $\sin x$ and $\cos x$
3. Add the Taylor series of $\cos \theta$ and $i \sin \theta$ together and you'll get the Taylor series for $e^{i\theta}$

Euler Identity and Polar-Cartesian Representations

In the previous section, we saw that a complex number $z = a + ib$ could be represented as a point (a, b) on the complex plane, which could also be viewed in polar coordinates as (r, θ) . We saw that to convert between the Cartesian (a, b) and the Polar (r, θ) representations, only simple trigonometry and Pythagoras is needed:

$a + ib \rightarrow r \angle \theta$	$r \angle \theta \rightarrow a + ib$
$r^2 = a^2 + b^2$	$a = r \cos \theta$
$\tan \theta = \frac{b}{a}$	$b = r \sin \theta$

This means that:

$$z = a + ib = r \cos \theta + i r \sin \theta = r (\cos \theta + i \sin \theta) = r e^{i\theta}$$

This offers another interpretation of the Euler identity as the algebraic conversion between Cartesian and Polar coordinates:

	Cartesian	Polar
Graphical	(a, b)	(r, θ)
Algebraic	$z = a + ib$	$z = r e^{i\theta}$

This now allows us to simplify a lot of difficult mathematics. For example let's look at the root problem $z^3 = 8$. Since the number 8 on the complex plane is the point $(8, 0)$, in polar coordinates, it can be any of the following: $8 \angle 0, 8 \angle 2\pi, 8 \angle 4\pi, \dots$ This is because we can go around the circle as many times as we want and return to the same point. Since we expect three roots, let's use the first three polar representations of 8:

$$z^3 = 8 = \left\{ 8e^{0i}, 8e^{2\pi i}, 8e^{4\pi i}, \dots \right\}$$

$$\begin{aligned} \rightarrow z &= \left\{ \left(8e^{0i} \right)^{\frac{1}{3}}, \left(8e^{2\pi i} \right)^{\frac{1}{3}}, \left(8e^{4\pi i} \right)^{\frac{1}{3}}, \dots \right\} \\ e^{\frac{0i}{3}} &= 2, \quad e^{\frac{2\pi i}{3}} = 2 \left(\cos \frac{2\pi}{3} + i \sin \frac{2\pi}{3} \right) = 2 \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) = -1 + i\sqrt{3} \\ e^{\frac{4\pi i}{3}} &= 2 \left(\cos \frac{4\pi}{3} + i \sin \frac{4\pi}{3} \right) = 2 \left(-\frac{1}{2} - i \frac{\sqrt{3}}{2} \right) = -1 - i\sqrt{3} \end{aligned}$$

If we had used more than 3 numbers, the roots would have started repeating. If we had used less than 3, we would have missed some answers in the same way that there are two answers to $z^2 = 9$ (namely $z = \pm 3$).

Cartesian vs Polar

Which is the best representation: Cartesian, $z = a + ib$, or Polar, $z = re^{i\theta}$. As you might expect, it depends on what you're trying to do... For example, let's take:

$$z_1 = 1 + i = \sqrt{2}e^{i\frac{\pi}{4}} \quad \text{and} \quad z_2 = -1 + i = \sqrt{2}e^{i\frac{3\pi}{4}}$$

Imagine having to add, subtract, multiply, divide or divide these together. Or raise them to a power, or take a root of them. Which of the two representations do you think would be easiest to use for each operation?

Operation	Cartesian	Polar
Addition	Easiest: $z_1 + z_2 = (1 + i) + (-1 + i) = 2i$	I don't know how to do it. $z_1 + z_2 = \sqrt{2}e^{i\frac{\pi}{4}} + \sqrt{2}e^{i\frac{3\pi}{4}} = ??$
Subtraction	Easiest: $z_1 - z_2 = (1 + i) - (-1 + i) = 2$	
Multiplication	Moderate: $z_1 \cdot z_2 = (1 + i) \cdot (-1 + i) = (1)(-1) + (1)(i) + (i)(-1) + (i)(i) = -1 + i - i - 1 = -2$	Easiest: $z_1 \cdot z_2 = \sqrt{2}e^{i\frac{\pi}{4}} \cdot \sqrt{2}e^{i\frac{3\pi}{4}} = \sqrt{2} \cdot \sqrt{2} e^{i(\frac{\pi}{4} + \frac{3\pi}{4})} = 2e^{i\pi} = -2$
Division	Tedious: $\frac{z_1}{z_2} = \frac{(1 + i)}{(-1 + i)} = \frac{(1 + i)(-1 - i)}{(-1 + i)(-1 - i)} = \frac{-1 - i + i - 1}{1 + 1} = \frac{-2 - i}{2} = -1 - \frac{i}{2}$	Easiest: $\frac{z_1}{z_2} = \frac{\sqrt{2}e^{i\frac{\pi}{4}}}{\sqrt{2}e^{i\frac{3\pi}{4}}} = e^{i(\frac{\pi}{4} - \frac{3\pi}{4})} = e^{-i\frac{\pi}{2}} = -i$
Exponentiation	The bigger the exponent, the more tedious: $z_1^{100} = (1 + i)^{100}$	Easy no matter how big the exponent: $z_1^{100} = (\sqrt{2}e^{i\frac{\pi}{4}})^{100} = 2^{50}e^{i25\pi} = 2^{50}e^{i(12\pi + \pi)} = 2^{50}(1 - i) = -2^{50}i$
Roots	I don't know if it's even possible: $\sqrt[3]{z_2} = \sqrt[3]{-1 + i}$	As easy as exponentiation: $\sqrt[3]{z_2} = (\sqrt{2}e^{i\frac{3\pi}{4}})^{\frac{1}{3}} = \sqrt[3]{2}e^{i\frac{\pi}{4}} = \frac{1 + i}{\sqrt{3}}$

The lesson here is that since the polar representation uses exponents, and exponents turn multiplication into addition⁴⁾, the polar representation is easiest for multiplication, division, exponentiation, and roots. It's essentially why the [dB scale](#) is so useful.

Important Algebraic Results

- Use the Euler identity to get the following two useful results:

$$\cos \theta = \frac{e^{i\theta} + e^{-i\theta}}{2} \quad \text{and} \quad \sin \theta = \frac{e^{i\theta} - e^{-i\theta}}{2i}$$

- Modify the Euler identity to see what happens when the angle is $(-\theta)$

$$\begin{aligned} e^{i\theta} &= \cos \theta + i \sin \theta \\ e^{-i\theta} &= \cos(-\theta) + i \sin(-\theta) \end{aligned}$$

- Add the original Euler identity to the new one for $(-\theta)$ and simplify:

$$\begin{aligned} e^{i\theta} + e^{-i\theta} &= \cos \theta + i \sin \theta + \cos \theta - i \sin \theta \\ &= 2 \cos \theta \end{aligned}$$

- To get the second equation, take the original Euler identity and subtract the new one for $(-\theta)$ from it and simplify:

$$\begin{aligned} e^{i\theta} - e^{-i\theta} &= \cos \theta + i \sin \theta - (\cos \theta - i \sin \theta) \\ &= 2i \sin \theta \end{aligned}$$

- Use the Euler identity to show that:

$$\begin{aligned} \cos(\theta + \phi) &= \cos \theta \cos \phi - \sin \theta \sin \phi \\ \sin(\theta + \phi) &= \sin \theta \cos \phi + \cos \theta \sin \phi \end{aligned}$$

- Multiply the Euler identities for θ and ϕ and simplify.

$$\begin{aligned} e^{i\theta} e^{i\phi} &= (\cos \theta + i \sin \theta)(\cos \phi + i \sin \phi) \\ &= \cos \theta \cos \phi - \sin \theta \sin \phi + i(\sin \theta \cos \phi + \cos \theta \sin \phi) \\ &= \cos(\theta + \phi) + i \sin(\theta + \phi) \end{aligned}$$

But the Euler identity for angle $(\theta + \phi)$ is: $e^{i(\theta + \phi)} = \cos(\theta + \phi) + i \sin(\theta + \phi)$

So comparing the **real** and **imaginary** parts, we can conclude that: $\cos(\theta + \phi) = \cos \theta \cos \phi - \sin \theta \sin \phi$ and $\sin(\theta + \phi) = \sin \theta \cos \phi + \cos \theta \sin \phi$

- Use the two earlier results to show that:

$$\sin(\theta + \Delta\theta) + \sin(\theta - \Delta\theta) = 2 \cos \Delta\theta \sin \theta$$

- Use the last result for $\sin(\theta + \phi)$ but replace ϕ with $\Delta\theta$

$$\sin(\theta + \Delta\theta) = \cos \theta \sin \Delta\theta + \sin \theta \cos \Delta\theta$$

Similarly, $\sin(\theta - \Delta\theta) = \cos \theta \sin(-\Delta\theta) + \sin \theta \cos(-\Delta\theta)$
 $\sin(\theta - \Delta\theta) = -\cos \theta \sin \Delta\theta + \sin \theta \cos \Delta\theta$

Adding both of these together, we get:
 $\sin(\theta + \Delta\theta) + \sin(\theta - \Delta\theta) = \cos \theta \sin \Delta\theta + \sin \theta \cos \Delta\theta - \cos \theta \sin \Delta\theta + \sin \theta \cos \Delta\theta = 2 \sin \theta \cos \Delta\theta$

This last result is the basis behind why [modulating the amplitude](#) of a carrier produces side bands.

Differential Equations

In the physics of wave, we often have to find solutions to the following type of differential equations:

$$a \ddot{x}(t) + b \dot{x}(t) + c x(t) = 0$$



A differential equation is an equation that relates a function to its derivatives in some ways and the question is: given some information about the system, what's the function (or family of functions) that satisfy the differential equation.

In physics we often use a dot above the function to indicate a derivative with respect to time, where as in math, we'll often use an apostrophe. Physicists don't like the apostrophe too much because they sometimes use it to denote a different coordinate system. So don't let the notation confuse you: $\dot{x}(t) = x'(t) = \frac{dx}{dt}$ and $\ddot{x}(t) = x''(t) = \frac{d^2x}{dt^2}$

In our applications, the parameters a, b, c are all real and positive quantities. Even without having studied differential equations in any depth, we can imagine that a possible solution to the above differential equation would be: $x(t) = e^{rt}$ since the derivative of an exponential function is itself an exponential function, which is encouraging.

The next step is to try this —“test function” in the differential equation and see if we can find the values of r that make it work. First we'll need derivatives of the test function:

$$\begin{aligned} & \text{\& } x(t) = e^{rt} \text{\& } \Rightarrow \text{\& } \dot{x}(t) = r e^{rt} \text{\& } \Rightarrow \text{\& } \ddot{x}(t) = r^2 e^{rt} \text{\& } \end{aligned}$$

When we put these into the differential equation, we get:

$$\begin{aligned} & a \ddot{x}(t) + b \dot{x}(t) + c x(t) = 0 \Rightarrow a (r^2 e^{rt}) + b (r e^{rt}) + c (e^{rt}) = 0 \Rightarrow e^{rt} (a r^2 + b r + c) = 0 \Rightarrow a r^2 + b r + c = 0 \Rightarrow \\ & r = -\frac{b}{2a} \pm \frac{\sqrt{b^2 - 4ac}}{2a} \end{aligned}$$

So what does that result mean? Remember, what we're looking for is the function $x(t)$ that satisfies the differential equation $a \ddot{x}(t) + b \dot{x}(t) + c x(t) = 0$

What we've got so far says that our test function $x(t) = e^{rt}$ will satisfy the differential equation if r is given by above equation. There is still a lot to unpack however. For example, since r contains a square root, it could be real or complex depending on the values of $a, b,$ and c . And as we saw above, if r is real, then $x(t)$ will be a real exponential function. But if r is complex, then we can expect $x(t)$ to be some sort of sinusoidal function (recall the Euler Identity).

To simplify the notation, let's define α and β as: $\alpha = \frac{b}{2a} \text{ and } \beta = \frac{\sqrt{b^2 - 4ac}}{2a}$

Notice how the absolute value under the square root ensures that β is always real.

r then:

$$r = \begin{cases} -\alpha \pm \beta & \text{if } b^2 - 4ac > 0, \\ -\alpha \pm i\beta & \text{if } b^2 - 4ac < 0, \end{cases}$$

Let's examine both of these cases in more detail.

Case 1: Over Damped Oscillation

When $b^2 - 4ac > 0$, r is real and the general solution is:

$$\begin{aligned} x(t) &= A_1 e^{r_1 t} + A_2 e^{r_2 t} \Rightarrow A_1 e^{(-\alpha + \beta) t} + A_2 e^{(-\alpha - \beta) t} \Rightarrow \\ &= A_1 e^{-\alpha t} e^{\beta t} + A_2 e^{-\alpha t} e^{-\beta t} \end{aligned}$$

$$x(t) = e^{-\alpha t} (A_1 e^{\beta t} + A_2 e^{-\beta t})$$

It's normal to have two constants of integration since our differential equation has a second degree derivative in it. To find these constants, we'd need to know more about the system's initial conditions.

Case 2: Under Damped

When $b^2 - 4ac < 0$, r is complex and we'll be using the Euler identity to simplify our solutions

$$\begin{aligned} x(t) &= A_1 e^{r_1 t} + A_2 e^{r_2 t} \Rightarrow A_1 e^{(-\alpha + i\beta) t} + A_2 e^{(-\alpha - i\beta) t} \Rightarrow \\ &= A_1 e^{-\alpha t} e^{i\beta t} + A_2 e^{-\alpha t} e^{-i\beta t} \Rightarrow e^{-\alpha t} (A_1 e^{i\beta t} + A_2 e^{-i\beta t}) \Rightarrow \\ &= e^{-\alpha t} \Big(A_1 (\cos(\beta t) + i \sin(\beta t)) + A_2 (\cos(\beta t) - i \sin(\beta t)) \Big) \Rightarrow \\ &= e^{-\alpha t} \Big((A_1 + A_2) \cos(\beta t) + i(A_1 - A_2) \sin(\beta t) \Big) \Rightarrow \\ &= e^{-\alpha t} (a_1 \cos(\beta t) + a_2 \sin(\beta t)) \end{aligned}$$

$$\cos(\beta t) + A \cos \phi \sin(\beta t) \Big) \quad \& \quad A e^{-\alpha t} \Big(\sin \phi \cos(\beta t) + \cos \phi \sin(\beta t) \Big) \quad \end{align*}$$

In the last three lines, we've redefined the constants of integration a few times so that:

$$\begin{align*} a_1 &= A_1 + A_2 \quad , \quad a_2 = i(A_1 - A_2) \quad \& \quad a_1 = A \sin \phi \quad , \quad a_2 = A \cos \phi \end{align*}$$

And we finally use one of the trig identities we proved earlier to write the solution as: $x(t) = A e^{-\alpha t} \sin(\beta t + \phi)$

Case 3: Critically Damped

When $b^2 - 4ac = 0$, $r = -\frac{b}{2a}$ is real and negative but our test solution is under determined. We'll instead propose a solution of the following type and test that it works:
$$\begin{align*} \&\& x(t) &= e^{rt}(A + Bt) \quad \&\& \dot{x}(t) = re^{rt}(A + Bt) + Be^{rt} \quad \&\& = e^{rt} \big(r(A + Bt) + B\big) \quad \&\& = e^{rt}(rA + B + Bt) \quad \&\& \ddot{x}(t) = re^{rt}(rA + B + Bt) + e^{rt}Br \quad \&\& = e^{rt} \big(r(rA + B + Bt) + Br\big) \quad \&\& = e^{rt}(Ar^2 + 2Br + Br^2t) \end{align*}$$

Example

Nous avons donc deux types de solutions complètement différents qui dépendent de trois paramètres a, b, c . Pour voir comment ces paramètres affectent le graphique, imaginons qu'une de nos conditions initiales est $\phi = \frac{\pi}{2}$. Ça veut dire que:

$$\begin{align*} \&\& a_1 = A \sin \pi/2 = A \quad \&\& a_2 = A \cos \pi/2 = 0 \quad \&\& \&\& A_1 + A_2 = A \quad \&\& A_1 - A_2 = 0 \quad \&\& \&\& A_1 = A/2 \quad \&\& A_2 = A/2 \end{align*}$$

Dans ce cas particulier, nous avons donc:

$$\begin{equation*} x(t) = \left\{ \begin{array}{l} A e^{-\alpha t} \frac{e^{\beta t} + e^{-\beta t}}{2} \quad \&\& b^2 - 4ac > 0, \\ A e^{-\alpha t} \cos(\beta t) \quad \&\& b^2 - 4ac < 0, \end{array} \right. \end{equation*}$$



Geogebra

1)

Whether math is discovered or invented is a famous philosophical problem. If you think it's invented: does that mean that $2 + 2$ didn't equal 4 until someone invented that? If you think it's discovered, what about computer programs? At the root, all computing is basically just math.

2)

Number line picture from [Wikipedia: Number line](#)

3)

Note, to obtain the [most beautiful equation in the world](#), set $\theta = \pi$ in the Euler identity:

$$e^{i\pi} = \cos(\pi) + i \sin(\pi) = -1 + 0i = -1 \quad \&\& \&\& e^{i\pi} = -1$$
 which is amazingly beautiful because it relates $e = 2.71828...$, $i = \sqrt{-1}$, $\pi = 3.14159...$, and -1 in the most surprising and elegant way.

4)

$$x^a \cdot x^b = x^{a+b}$$